

Foreword to the Special Issue on Spectral Unmixing of Remotely Sensed Data

MORE than two decades after the first efforts toward the application of spectral mixture analysis techniques to remotely sensed data [1], [2], effective spectral unmixing still remains an elusive exploitation goal. Regardless of the available spatial resolution, the spectral signals collected in natural environments are invariably a mixture of the signatures of the various materials found within the spatial extent of the ground instantaneous field view of the remote sensing imaging instrument [3]. The availability of hyperspectral imaging instruments [4] (also called imaging spectrometers [5]) with a number of spectral bands that exceeds the number of spectral mixture components has fostered many research efforts. Spectral unmixing has been a very active research area in recent years since it faces important challenges [6], [7].

In order to present the state-of-the-art and most recent developments in this area, it is our great pleasure to introduce this special issue on *Spectral Unmixing of Remotely Sensed Data* of the IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING. The special issue is the first one of its kind in the literature since this topic has not been addressed in the form of a dedicated monograph in any other journal in the past. The special issue brings together distinguished experts to provide a remarkable sample of latest-generation techniques in the field. A large number of submissions (45) were received for this special issue, of which 20 papers were selected after rigorous review. In the remainder of this foreword, we review key issues and topics of current interest related to spectral unmixing that are covered by this special issue.

A. Linear and Nonlinear Spectral Unmixing

Linear spectral unmixing [8] is a standard technique for spectral mixture analysis that infers a set of pure spectral signatures, called *endmembers* [9], [10], and the fractions of these endmembers, called *abundances* [11], in each pixel of the scene. This model assumes that the spectra collected by the imaging spectrometer can be expressed in the form of a linear combination of endmembers, weighted by their corresponding abundances. Because each observed spectral signal is the result of an actual mixing process, it is expected that the driving abundances satisfy two constraints, i.e., they should be nonnegative [12] and the sum of abundances for a given pixel should be unity [13]. Although the linear model has practical advantages, such as ease of implementation and flexibility in different

applications, nonlinear unmixing describes mixed spectra (in physical [14], [15], or statistical [16] sense) by assuming that part of the source radiation is multiply scattered before being collected at the sensor. The distinction between the linear and the nonlinear unmixing has been widely studied in recent years [17].

In this special issue, several contributions are directly related to these topics. In [18], improvements over fully constrained least squares techniques for linear spectral unmixing [11] are discussed. In [19], a new component analysis-based strategy for linear spectral unmixing is presented. These techniques assume that all endmember signatures are available in advance. In the case that this requirement is not satisfied, partial unmixing has emerged as a suitable alternative to solve the abundance estimation problem. One widely used partial unmixing technique is mixture-tuned matched filtering (MTMF), which is described in detail in another contribution of this special issue [20]. Additional results of MTMF are also illustrated for the Cuprite Mining District, Nevada, USA, on this special issue cover.

Finally, two new techniques for nonlinear spectral unmixing are also introduced in this special issue. In [21], a generalized bilinear model and a hierarchical Bayesian algorithm for nonlinear unmixing of hyperspectral images are proposed, following previous efforts on the application of Bayesian techniques to spectral unmixing [22], [23]. In [24], a novel approach based on neural networks for the extraction of pixel abundances from hyperspectral data is developed. This work expands over previous efforts in the literature focused on using neural networks for nonlinear unmixing purposes [25]–[28].

B. Endmember Determination and Pure Class Modeling

Early approaches to endmember determination were principally manual [29], [30]. More recent development of automatic or semiautomatic endmember extraction algorithms has resulted in significant steps forward. While many available approaches associate a single spectral signature to each endmember, multiple endmembers have also been used to account for within (pure) class spectral variation [31]–[34]. When multiple spectra are used to represent a pure class, the term endmember refers to all spectra in the modeled pure class.

Notwithstanding the importance of multiple endmember-based approaches, a majority of algorithms have been designed under the pure pixel assumption, i.e., they assume that the remotely sensed data contain one pure observation for each distinct material present in the scene. This allows validation of extracted endmembers with regard to reference signatures

using different distance metrics [35]. Perhaps, due to their ease of computation and clear conceptual meaning, these are the most widely used class of algorithms for endmember determination, with a plethora of algorithms designed under this assumption (see [36]–[61], among several others). It should be noted, however, that some of the aforementioned algorithms require a dimensionality estimation step [62]. Under the linear mixture assumption, this amounts to estimating the number of endmembers [63], [64].

This special issue includes two new contributions on endmember determination under the pure pixel assumption. In [65], a new algorithm is developed for this purpose based on discrete particle swarm optimization concepts. In [66], the well-known N-FINDR endmember extraction algorithm originally developed in [41] is revisited, and two new algorithms are derived from alternative optimization strategies, namely, alternating optimization and successive optimization.

C. Endmember Determination Without Pure Pixel Assumption

Although the maximum volume procedure adopted by N-FINDR and related algorithms is successful when pure signatures are present in the data, given the available spatial resolution of state-of-the-art imaging spectrometers and the presence of the mixture phenomenon at different scales (even at microscopic levels), in some cases, the pure pixel assumption may not be valid. To address this issue, several endmember determination techniques have been developed without assuming the presence of pure signatures in the input data. These methods aim at generating *virtual* endmembers [67] (not necessarily present in the set comprised by input data samples) by finding the simplex with minimum volume that encompasses all observations [68]–[73].

In this special issue, a new minimum-volume enclosing algorithm is developed [74]. It accounts for the noise effects in the observations by employing chance constraints and improves over a previously developed algorithm [72].

D. Incorporation of Spatial Information Into Endmember Determination and Spectral Unmixing

Most of the techniques discussed so far for endmember determination and spectral unmixing rely on the exploitation of spectral information alone. However, one of the distinguishing properties of remotely sensed data is the multivariate information coupled with a 2-D (pictorial) representation amenable to image interpretation. Subsequently, unmixing techniques can benefit from an integrated framework in which both the spectral information and the spatial arrangement of pixel vectors are taken into account. This aspect has been widely studied in the unmixing literature [75]–[81].

In this special issue, four contributions are directly related with this issue, including the development of new techniques for spatial–spectral endmember determination [82], [83] and the incorporation of spatial information into hyperspectral image unmixing [84], [85].

E. Sparse Regression-Based Unmixing

A recently developed approach to tackle the problems related to the unavailability of pure spectral signatures is to model mixed pixel observations as linear combinations of spectra from a library collected on the ground by a field spectroradiometer. Unmixing then amounts to finding the optimal subset of signatures in a (potentially very large) spectral library that can best model each mixed pixel in the scene [86]. In practice, this is a combinatorial problem that calls for efficient sparse regression techniques based on sparsity-inducing regularizers since the number of endmembers participating in a mixed pixel is usually very small compared with the (ever-growing) dimensionality and availability of spectral libraries.

In this special issue, two contributions are provided on the topic of sparse regression applied to spectral unmixing. In [87], a new method for subpixel modeling, mapping, and classification of hyperspectral images is presented. It uses learned block-structured discriminative dictionaries [88], where each block is adapted and optimized to represent a specific material in a compact and sparse manner. A spatial–spectral coherence regularizer is also applied. In [89], a sparsity constraint is included in nonnegative matrix factorization, a widely used technique to unmix hyperspectral images and recover the material endmembers [90], [91].

F. Unmixing of Remotely Sensed Data With Moderate Spectral Resolution

Although most of the techniques discussed so far have been designed to unmix remotely sensed hyperspectral scenes with a large number of spectral bands, such as those provided by the National Aeronautics and Space Administration (NASA) Airborne Visible/Infrared Imaging Spectrometer [92] or the Hyperion instrument onboard the Earth Observing 1 satellite [93], the unmixing of scenes with moderate spectral resolution is also of high interest, as high spectral resolution data are not always available.

In this special issue, two contributions deal with this relevant issue. In [94], a new method to unmix remotely sensed data collected by NASA's moderate resolution imaging spectrometer is presented. In [95], a time series of images collected by the European Space Agency's Medium Resolution Imaging Spectrometer is unmixed, both for each image and also by layer stacking all the images in the time series, thus addressing multitemporal spectral unmixing problems.

G. Connections Between Spectral Unmixing and Classification

Spectral unmixing and *hard* classification methods [96], [97] can be seen as complementary techniques since the latter are more suitable for the classification of pixels dominated by a single land cover class, while the former are devoted to the analysis of mixed pixels and can be seen as a form of *soft* classification. Because remotely sensed images contain areas with both pure and mixed pixels, the combination of these two techniques provides an interesting analysis approach.

This topic has been explored in previous contributions [33], [98]–[100].

In this special issue, the contribution [101] addresses this synergistic data processing trend by developing a new unmixing-to-classification conversion model that treats the abundance quantification task as a classification problem.

H. Applications of Spectral Unmixing

Numerous applications related to the monitoring of the environment and the retrieval of biogeophysical parameters have been addressed using spectral unmixing techniques, covering (to name a few) vegetative [102]–[104], soil [105]–[107], urban [108]–[110], and planetary [111]–[113] surfaces. Three application-oriented contributions are included in this special issue. In [114], the retrieval of canopy closure and leaf area index from Moso bamboo (an important forest type in subtropical areas of China) is investigated through linear spectral unmixing. In [115], the capability of spectral unmixing for analyzing hyperspectral images from Mars is explored. In [116], the unmixing of atmospheric trace gases from hyperspectral satellite data is addressed.

Combined, the different topics included in this special issue provide an excellent snapshot of the state of the art in the area of spectral unmixing, offering a thoughtful perspective on the potential and emerging challenges of applying unmixing techniques to different types of remotely sensed data. The Guest Editors would like to take this opportunity to gratefully thank the Editor-in-Chief, Prof. Christopher S. Ruf, for his constant support and encouragement to this special issue. The Guest Editors also gratefully thank all the contributors and reviewers who participated in the evaluation of manuscripts for the special issue. Without their outstanding contributions, the special issue could not have been completed.

ANTONIO PLAZA, *Guest Editor*
Department of Technology of Computers and
Communications
University of Extremadura
10003 Caceres, Spain

QIAN DU, *Guest Editor*
Department of Electrical and Computer
Engineering
Mississippi State University
Mississippi State, MS 39762 USA

JOSÉ M. BIOCAS-DIAS, *Guest Editor*
Department of Electrical and Computer
Engineering
Instituto Superior Tecnico
1049-001 Lisbon, Portugal

XIUPING JIA, *Guest Editor*
Department of Electrical Engineering
School of Engineering and Information
Technology
University College
University of North South Wales
2904 Canberra, Australia

FRED A. KRUSE, *Guest Editor*
Physics Department
Naval Postgraduate School
Monterey, CA 93943 USA

REFERENCES

- [1] P. E. Johnson, M. O. Smith, S. Taylor-George, and J. B. Adams, "A semiempirical method for analysis of the reflectance spectra for binary mineral mixtures," *J. Geophys. Res.*, vol. 88, no. B4, pp. 3557–3561, 1983.
- [2] J. B. Adams, M. O. Smith, and P. E. Johnson, "Spectral mixture modeling: A new analysis of rock and soil types at the Viking Lander 1 site," *J. Geophys. Res.*, vol. 91, no. B8, pp. 8098–8112, 1986.
- [3] N. Keshava and J. F. Mustard, "Spectral unmixing," *IEEE Signal Process. Mag.*, vol. 19, no. 1, pp. 44–57, Jan. 2002.
- [4] A. Plaza, J. A. Benediktsson, J. Boardman, J. Brazile, L. Bruzzone, G. Camps-Valls, J. Chanussot, M. Fauvel, P. Gamba, J. Gualtieri, M. Marconcini, J. C. Tilton, and G. Trianni, "Recent advances in techniques for hyperspectral image processing," *Remote Sens. Environ.*, vol. 113, no. Supplement 1, pp. 110–122, Sep. 2009.
- [5] M. E. Schaepman, S. L. Ustin, A. Plaza, T. H. Painter, J. Verrelst, and S. Liang, "Earth system science related imaging spectroscopy—An assessment," *Remote Sens. Environ.*, vol. 113, pp. 123–137, 2009.
- [6] J. M. Bioucas-Dias and A. Plaza, "Hyperspectral unmixing: Geometrical, statistical, and sparse regression-based approaches," in *Proc. SPIE—Image and Signal Processing for Remote Sensing XVI*, Toulouse, France, Sep. 20–23, 2010, vol. 7830, pp. 1–15.
- [7] M. Parente and A. Plaza, "Survey of geometric and statistical unmixing algorithms for hyperspectral images," in *Proc. 2nd IEEE WHISPERS—Evolution in Remote Sensing*, Reykjavik, Iceland, Jun. 14–16, 2010, pp. 1–4.
- [8] J. J. Settle and N. A. Drake, "Linear mixing and the estimation of ground cover proportions," *Int. J. Remote Sens.*, vol. 14, no. 6, pp. 1159–1177, 1993.
- [9] A. Plaza, P. Martinez, R. Perez, and J. Plaza, "A quantitative and comparative analysis of endmember extraction algorithms from hyperspectral data," *IEEE Trans. Geosci. Remote Sens.*, vol. 42, no. 3, pp. 650–663, Mar. 2004.
- [10] Q. Du, N. Raksuntorn, N. H. Younan, and R. L. King, "End-member extraction for hyperspectral image analysis," *Appl. Opt.*, vol. 47, no. 28, pp. 77–84, Oct. 2008.
- [11] D. Heinz and C.-I. Chang, "Fully constrained least squares linear mixture analysis for material quantification in hyperspectral imagery," *IEEE Trans. Geosci. Remote Sens.*, vol. 39, no. 3, pp. 529–545, Mar. 2001.
- [12] C.-I. Chang and D. Heinz, "Constrained subpixel target detection for remotely sensed imagery," *IEEE Trans. Geosci. Remote Sens.*, vol. 38, no. 3, pp. 1144–1159, May 2000.
- [13] C.-I. Chang, *Hyperspectral Imaging: Techniques for Spectral Detection and Classification*. New York: Plenum, 2003.
- [14] C. C. Borel and S. A. W. Gerstl, "Nonlinear spectral mixing model for vegetative and soil surfaces," *Remote Sens. Environ.*, vol. 47, no. 3, pp. 403–416, Mar. 1994.
- [15] W. Liu and E. Y. Wu, "Comparison of non-linear mixture models," *Remote Sens. Environ.*, vol. 94, no. 2, pp. 145–154, Jan. 2005.
- [16] N. Raksuntorn and Q. Du, "Nonlinear spectral mixture analysis for hyperspectral imagery in an unknown environment," *IEEE Geosci. Remote Sens. Lett.*, vol. 7, no. 4, pp. 836–840, Oct. 2010.
- [17] A. Plaza, G. Martín, J. Plaza, M. Zortea, and S. Sánchez, "Recent developments in spectral unmixing and endmember extraction," in *Opt. Remote Sens.*, S. Prasad, L. M. Bruce, and J. Chanussot, Eds. Berlin, Germany: Springer-Verlag, 2011, ch. 12, pp. 235–267.
- [18] R. Heylen, D. Burazerović, and P. Scheunders, "Fully constrained least squares spectral unmixing by simplex projection," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4112–4122, Nov. 2011.
- [19] C.-I. Chang, X. Jiao, C.-C. Wu, E. Y. Du, and H.-M. Chen, "Component analysis-based unsupervised linear spectral mixture analysis for hyperspectral imagery," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4123–4137, Nov. 2011.
- [20] J. W. Boardman and F. A. Kruse, "Analysis of imaging spectrometer data using N-dimensional geometry and a mixture-tuned matched filtering (MTMF) approach," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4138–4152, Nov. 2011.

- [21] A. Halimi, Y. Altmann, N. Dobigeon, and J.-Y. Tourneret, "Nonlinear unmixing of hyperspectral images using a generalized bilinear model," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4153–4162, Nov. 2011.
- [22] N. Dobigeon, S. Moussaoui, M. Coulon, J.-Y. Tourneret, and A. O. Hero, "Joint Bayesian endmember extraction and linear unmixing for hyperspectral imagery," *IEEE Trans. Signal Process.*, vol. 57, no. 11, pp. 4355–4368, Nov. 2009.
- [23] O. Eches, N. Dobigeon, C. Mailhes, and J.-Y. Tourneret, "Bayesian estimation of linear mixtures using the normal compositional model application to hyperspectral imagery," *IEEE Trans. Image Process.*, vol. 19, no. 6, pp. 1403–1413, Jun. 2010.
- [24] G. A. Licciardi and F. Del Frate, "Pixel unmixing in hyperspectral data by means of neural networks," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4163–4172, Nov. 2011.
- [25] A. Baraldi, E. Binaghi, P. Blonda, P. A. Brivio, and P. Rampini, "Comparison of the multilayer perceptron with neuro-fuzzy techniques in the estimation of cover class mixture in remotely sensed data," *IEEE Trans. Geosci. Remote Sens.*, vol. 39, no. 5, pp. 994–1005, May 2001.
- [26] K. J. Guilfoyle, M. L. Althouse, and C.-I. Chang, "A quantitative and comparative analysis of linear and nonlinear spectral mixture models using radial basis function neural networks," *IEEE Trans. Geosci. Remote Sens.*, vol. 39, no. 10, pp. 2314–2318, Oct. 2001.
- [27] J. Plaza, A. Plaza, R. Perez, and P. Martinez, "On the use of small training sets for neural network-based characterization of mixed pixels in remotely sensed hyperspectral images," *Pattern Recognit.*, vol. 42, no. 11, pp. 3032–3045, Nov. 2009.
- [28] J. Plaza and A. Plaza, "Spectral mixture analysis of hyperspectral scenes using intelligently selected training samples," *IEEE Geosci. Remote Sens. Lett.*, vol. 7, no. 2, pp. 371–375, Apr. 2010.
- [29] C. A. Bateson and B. Curtiss, "A tool for manual endmember selection and spectral unmixing," in *Summaries 4th Annu. JPL Airborne Geoscience Workshop*, 1993, vol. 1, pp. 3–6.
- [30] C. A. Bateson and B. Curtiss, "A method for manual endmember selection and spectral unmixing," *Remote Sens. Environ.*, vol. 55, no. 3, pp. 229–243, Mar. 1996.
- [31] P. Dennison and D. Roberts, "Endmember selection for multiple endmember spectral mixture analysis using endmember average rmse," *Remote Sens. Environ.*, vol. 87, no. 2/3, pp. 123–135, Oct. 2003.
- [32] J. Settle, "On the effect of variable endmember spectra in the linear mixture model," *IEEE Trans. Geosci. Remote Sens.*, vol. 44, no. 2, pp. 389–396, Feb. 2006.
- [33] L. Wang and X. Jia, "Integration of soft and hard classifications using extended support vector machines," *IEEE Geosci. Remote Sens. Lett.*, vol. 6, no. 3, pp. 543–547, Jul. 2009.
- [34] B. Somers, G. P. Asner, L. Tits, and P. Coppin, "Endmember variability in spectral mixture analysis: A review," *Remote Sens. Environ.*, vol. 115, no. 7, pp. 1603–1616, Jul. 2011.
- [35] J. Chen, X. Jia, W. Yang, and B. Matsushita, "Generalization of subpixel analysis for hyperspectral data with flexibility in spectral similarity measures," *IEEE Trans. Geosci. Remote Sens.*, vol. 47, no. 7, pp. 2165–2171, Jul. 2009.
- [36] J. C. Harsanyi and C.-I. Chang, "Hyperspectral image classification and dimensionality reduction: An orthogonal subspace projection," *IEEE Trans. Geosci. Remote Sens.*, vol. 32, no. 4, pp. 779–785, Jul. 1994.
- [37] J. W. Boardman, F. A. Kruse, and R. O. Green, "Mapping target signatures via partial unmixing of AVIRIS data," in *Proc. JPL Airborne Earth Sci. Workshop*, 1995, pp. 23–26.
- [38] J. H. Bowles, P. J. Palmadesso, J. A. Antoniadis, M. M. Baumbach, and L. J. Rickard, "Use of filter vectors in hyperspectral data analysis," in *Proc. SPIE—Infrared Spaceborne Remote Sensing III*, 1995, vol. 2553, pp. 148–157.
- [39] R. A. Neville, K. Staenz, T. Szeredi, J. Lefebvre, and P. Hauff, "Automatic endmember extraction from hyperspectral data for mineral exploration," in *Proc. 21st Can. Symp. Remote Sens.*, 1999, pp. 21–24.
- [40] A. Ifarraguerri and C.-I. Chang, "Multispectral and hyperspectral image analysis with convex cones," *IEEE Trans. Geosci. Remote Sens.*, vol. 37, no. 2, pp. 756–770, Mar. 1999.
- [41] M. E. Winter, "N-FINDR: An algorithm for fast autonomous spectral endmember determination in hyperspectral data," in *Proc. SPIE*, 1999, vol. 3753, pp. 266–275.
- [42] Q. Du, H. Ren, and C.-I. Chang, "A comparative study for orthogonal subspace projection and constrained energy minimization," *IEEE Trans. Geosci. Remote Sens.*, vol. 41, no. 6, pp. 1525–1529, Jun. 2003.
- [43] M. Berman, H. Kiiveri, R. Lagerstrom, A. Ernst, R. Dunne, and J. F. Huntington, "ICE: A statistical approach to identifying endmembers in hyperspectral images," *IEEE Trans. Geosci. Remote Sens.*, vol. 42, no. 10, pp. 2085–2095, Oct. 2004.
- [44] J. M. P. Nascimento and J. M. Bioucas-Dias, "Vertex component analysis: A fast algorithm to unmix hyperspectral data," *IEEE Trans. Geosci. Remote Sens.*, vol. 43, no. 4, pp. 898–910, Apr. 2005.
- [45] A. Plaza and C.-I. Chang, "Impact of initialization on design of endmember extraction algorithms," *IEEE Trans. Geosci. Remote Sens.*, vol. 44, no. 11, pp. 3397–3407, Nov. 2006.
- [46] C.-I. Chang and A. Plaza, "A fast iterative algorithm for implementation of pixel purity index," *IEEE Geosci. Remote Sens. Lett.*, vol. 3, no. 1, pp. 63–67, Jan. 2006.
- [47] D. M. Rogge, B. Rivard, J. Zhang, and J. Feng, "Iterative spectral unmixing for optimizing per-pixel endmember sets," *IEEE Trans. Geosci. Remote Sens.*, vol. 44, no. 12, pp. 3725–3736, Dec. 2006.
- [48] J. Wang and C.-I. Chang, "Applications of independent component analysis in endmember extraction and abundance quantification for hyperspectral imagery," *IEEE Trans. Geosci. Remote Sens.*, vol. 44, no. 9, pp. 2601–2616, Sep. 2006.
- [49] C.-I. Chang, C.-C. Wu, W. Liu, and Y.-C. Ouyang, "A new growing method for simplex-based endmember extraction algorithm," *IEEE Trans. Geosci. Remote Sens.*, vol. 44, no. 10, pp. 2804–2819, Oct. 2006.
- [50] A. Zare and P. Gader, "Sparsity promoting iterated constrained endmember detection for hyperspectral imagery," *IEEE Geosci. Remote Sens. Lett.*, vol. 4, no. 3, pp. 446–450, Jul. 2007.
- [51] A. Zare and P. Gader, "Hyperspectral band selection and endmember detection using sparsity promoting priors," *IEEE Geosci. Remote Sens. Lett.*, vol. 5, no. 2, pp. 256–260, Apr. 2008.
- [52] M. Zortea and A. Plaza, "A quantitative and comparative analysis of different implementations of N-FINDR: A fast endmember extraction algorithm," *IEEE Geosci. Remote Sens. Lett.*, vol. 6, no. 4, pp. 787–791, Oct. 2009.
- [53] X. Tao, B. Wang, and L. Zhang, "Orthogonal bases approach for the decomposition of mixed pixels in hyperspectral imagery," *IEEE Geosci. Remote Sens. Lett.*, vol. 6, no. 2, pp. 219–223, Apr. 2009.
- [54] A. Zare and P. Gader, "PCE: Piecewise convex endmember detection," *IEEE Trans. Geosci. Remote Sens.*, vol. 48, no. 6, pp. 2620–2632, Jun. 2010.
- [55] C.-I. Chang, C.-C. Wu, C.-S. Lo, and M.-L. Chang, "Real-time simplex growing algorithms for hyperspectral endmember extraction," *IEEE Trans. Geoscience and Remote Sensing*, vol. 48, no. 4, pp. 1834–1850, Apr. 2010.
- [56] F. Schmidt, A. Schmidt, E. Tréguier, M. Guiheneuf, S. Moussaoui, and N. Dobigeon, "Implementation strategies for hyperspectral unmixing using Bayesian source separation," *IEEE Trans. Geosci. Remote Sens.*, vol. 48, no. 11, pp. 4003–4013, Nov. 2010.
- [57] O. Duran and M. Petrou, "Robust endmember extraction in the presence of anomalies," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 6, pp. 1986–1996, Jun. 2011.
- [58] B. Zhang, X. Sun, L. Gao, and L. Yang, "Endmember extraction of hyperspectral remote sensing images based on the ant colony optimization (ACO) algorithm," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 7, pp. 2635–2646, Jul. 2011.
- [59] M. Shoshany, F. Kizel, N. Netanyahu, N. Goldshlager, T. Jarmer, and G. Even-Tzur, "An iterative search in end-member fraction space for spectral unmixing," *IEEE Geosci. Remote Sens. Lett.*, vol. 8, no. 4, pp. 706–709, Jul. 2011.
- [60] C.-I. Chang, C.-C. Wu, and H.-M. Chen, "Random pixel purity index," *IEEE Geosci. Remote Sens. Lett.*, vol. 7, no. 2, pp. 324–328, Apr. 2010.
- [61] S. Dowler and M. Andrews, "On the convergence of N-FINDR and related algorithms: To iterate or not to iterate?," *IEEE Geosci. Remote Sens. Lett.*, vol. 8, no. 1, pp. 4–8, Jan. 2011.
- [62] A. A. Green, M. Berman, P. Switzer, and M. D. Craig, "A transformation for ordering multispectral data in terms of image quality with implications for noise removal," *IEEE Trans. Geosci. Remote Sens.*, vol. 26, no. 1, pp. 65–74, Jan. 1988.
- [63] C.-I. Chang and Q. Du, "Estimation of number of spectrally distinct signal sources in hyperspectral imagery," *IEEE Trans. Geosci. Remote Sens.*, vol. 42, no. 3, pp. 608–619, Mar. 2004.
- [64] J. M. Bioucas-Dias and J. M. P. Nascimento, "Hyperspectral subspace identification," *IEEE Trans. Geosci. Remote Sens.*, vol. 46, no. 8, pp. 2435–2445, Aug. 2008.
- [65] B. Zhang, X. Sun, L. Gao, and L. Yang, "Endmember extraction of hyperspectral remote sensing images based on the discrete particle swarm optimization algorithm," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4173–4176, Nov. 2011.

- [66] T.-H. Chan, W.-K. Ma, A. Ambikapathi, and C.-Y. Chi, "A simplex volume maximization framework for hyperspectral endmember extraction," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4177–4193, Nov. 2011.
- [67] X. Chen, J. Chen, X. Jia, B. Somers, J. Wu, and P. Coppin, "A quantitative analysis of virtual endmembers' increased impact on the collinearity effect in spectral unmixing," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 8, pp. 2945–2956, Aug. 2011.
- [68] M. D. Craig, "Minimum-volume transforms for remotely sensed data," *IEEE Trans. Geosci. Remote Sens.*, vol. 32, no. 3, pp. 542–552, May 1994.
- [69] L. Miao and H. Qi, "Endmember extraction from highly mixed data using minimum volume constrained nonnegative matrix factorization," *IEEE Trans. Geosci. Remote Sens.*, vol. 45, no. 3, pp. 765–777, Mar. 2007.
- [70] J. Li and J. Bioucas-Dias, "Minimum volume simplex analysis: A fast algorithm to unmix hyperspectral data," in *Proc. IEEE Int. Geosci. Remote Sens. Symp.*, 2008, vol. 3, pp. III-250–III-253.
- [71] J. Bioucas-Dias, "A variable splitting augmented Lagrangian approach to linear spectral unmixing," in *Proc. 1st IEEE WHISPER*, Grenoble, France, Aug. 26–28, 2009, pp. 1–4.
- [72] T.-H. Chan, C.-Y. Chi, Y.-M. Huang, and W.-K. Ma, "A convex analysis-based minimum-volume enclosing simplex algorithm for hyperspectral unmixing," *IEEE Trans. Signal Process.*, vol. 57, no. 11, pp. 4418–4432, Nov. 2009.
- [73] J. M. Bioucas-Dias and J. Nascimento, "Hyperspectral unmixing based on mixtures of Dirichlet components," *IEEE Trans. Geosci. Remote Sens.*, vol. 50, no. 3, Mar. 2012, to be published. [Online]. Available: DOI:10.1109/TGRS.2011.2163941
- [74] A. Ambikapathi, T.-H. Chan, W.-K. Ma, and C.-Y. Chi, "Chance-constrained robust minimum-volume enclosing simplex algorithm for hyperspectral unmixing," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4194–4209, Nov. 2011.
- [75] A. Plaza, P. Martinez, R. Perez, and J. Plaza, "Spatial/spectral endmember extraction by multidimensional morphological operations," *IEEE Trans. Geosci. Remote Sens.*, vol. 40, no. 9, pp. 2025–2041, Sep. 2002.
- [76] D. M. Rogge, B. Rivard, J. Zhang, A. Sanchez, J. Harris, and J. Feng, "Integration of spatial-spectral information for the improved extraction of endmembers," *Remote Sens. Environ.*, vol. 110, no. 3, pp. 287–303, 2007.
- [77] S. Jia and Y. Qian, "Spectral and spatial complexity-based hyperspectral unmixing," *IEEE Trans. Geosci. Remote Sens.*, vol. 45, no. 12, pp. 3867–3879, Dec. 2007.
- [78] M. Zortea and A. Plaza, "Spatial preprocessing for endmember extraction," *IEEE Trans. Geosci. Remote Sens.*, vol. 47, no. 8, pp. 2679–2693, Aug. 2009.
- [79] S. Mei, M. He, Z. Wang, and D. Feng, "Spatial purity based endmember extraction for spectral mixture analysis," *IEEE Trans. Geosci. Remote Sens.*, vol. 48, no. 9, pp. 3434–3445, Sep. 2010.
- [80] D. Thompson, L. Mandrake, M. Gilmore, and R. Castaño, "Superpixel endmember detection," *IEEE Trans. Geosci. Remote Sens.*, vol. 48, no. 11, pp. 4023–4033, Nov. 2010.
- [81] G. Martin and A. Plaza, "Region-based spatial preprocessing for endmember extraction and spectral unmixing," *IEEE Geosci. Remote Sens. Lett.*, vol. 8, no. 4, pp. 745–749, Jul. 2011.
- [82] S. Mei, M. He, Y. Zhang, Z. Wang, and D. Feng, "Improving spatial-spectral endmember extraction in the presence of anomalous ground objects," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4210–4222, Nov. 2011.
- [83] H. Li and L. Zhang, "A hybrid automatic endmember extraction algorithm based on a local window," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4223–4238, Nov. 2011.
- [84] O. Eches, N. Dobigeon, and J.-Y. Tourneret, "Enhancing hyperspectral image unmixing with spatial correlations," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4239–4247, Nov. 2011.
- [85] K. Canham, A. Schlamm, A. Ziemann, B. Basener, and D. Messinger, "Spatially adaptive hyperspectral unmixing," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4248–4262, Nov. 2011.
- [86] M. D. Iordache, J. Bioucas-Dias, and A. Plaza, "Sparse unmixing of hyperspectral data," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 6, pp. 2014–2039, Jun. 2011.
- [87] A. Castrodad, Z. Xing, J. B. Greer, E. Bosch, L. Carin, and G. Sapiro, "Learning discriminative sparse representations for modeling, source separation, and mapping of hyperspectral imagery," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4263–4281, Nov. 2011.
- [88] A. S. Charles, B. A. Olshausen, and C. J. Rozell, "Learning sparse codes for hyperspectral imagery," *IEEE J. Sel. Topics Appl. Earth Observations Remote Sens.*, vol. 5, no. 5, pp. 963–978, Sep. 2011.
- [89] Y. Qian, S. Jia, J. Zhou, and A. Robles-Kelly, "Hyperspectral unmixing via $l_{1/2}$ sparsity-constrained nonnegative matrix factorization," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4282–4297, Nov. 2011.
- [90] A. Huck, M. Guillaume, and J. Blanc-Talon, "Minimum dispersion constrained nonnegative matrix factorization to unmix hyperspectral data," *IEEE Trans. Geosci. Remote Sens.*, vol. 48, no. 6, pp. 2590–2602, Jun. 2010.
- [91] X. Liu, W. Xia, B. Wang, and L. Zhang, "An approach based on constrained nonnegative matrix factorization to unmix hyperspectral data," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 2, pp. 757–772, Feb. 2011.
- [92] R. O. Green, M. L. Eastwood, C. M. Sarture, T. G. Chrien, M. Aronsson, B. J. Chippendale, J. A. Faust, B. E. Pavri, C. J. Chovit, M. Solis, M. R. Olah, and O. Williams, "Imaging spectroscopy and the airborne visible/infrared imaging spectrometer (AVIRIS)," *Remote Sens. Environ.*, vol. 65, no. 3, pp. 227–248, Sep. 1998.
- [93] F. A. Kruse, J. W. Boardman, and J. F. Huntington, "Comparison of airborne hyperspectral data and EO-1 Hyperion for mineral mapping," *IEEE Trans. Geosci. Remote Sens.*, vol. 41, no. 6, pp. 1388–1400, Jun. 2003.
- [94] H. Li, Y. Wang, Y. Li, and X. Wang, "Pixel-unmixing moderate-resolution remote sensing imagery using pairwise coupling support vector machines: A case study," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4298–4307, Nov. 2011.
- [95] R. Zurita-Milla, L. Gómez-Chova, L. Guanter, J. G. P. W. Clevers, and G. Camps-Valls, "Multitemporal unmixing of medium-spatial-resolution satellite images: A case study using MERIS images for land-cover mapping," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4308–4317, Nov. 2011.
- [96] D. A. Landgrebe, *Signal Theory Methods in Multispectral Remote Sensing*. New York: Wiley, 2003.
- [97] J. A. Richards and X. Jia, *Remote Sensing Digital Image Analysis: An Introduction*. New York: Springer-Verlag, 2006.
- [98] A. Plaza, P. Martinez, R. M. Perez, and J. Plaza, "A new approach to mixed pixel classification of hyperspectral imagery based on extended morphological profiles," *Pattern Recognit.*, vol. 37, no. 6, pp. 1097–1116, 2004.
- [99] I. Dopido, M. Zortea, A. Villa, A. Plaza, and P. Gamba, "Unmixing prior to supervised classification of remotely sensed hyperspectral images," *IEEE Geosci. Remote Sens. Lett.*, vol. 8, no. 4, pp. 760–764, Jul. 2011.
- [100] A. Villa, J. Chanussot, J. A. Benediktsson, and C. Jutten, "Spectral unmixing for the classification of hyperspectral images at a finer spatial resolution," *IEEE J. Sel. Topics Signal Process.*, vol. 5, no. 3, pp. 521–533, Jun. 2011.
- [101] F. A. Mianji and Y. Zhang, "SVM-based unmixing-to-classification conversion for hyperspectral abundance quantification," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4318–4327, Nov. 2011.
- [102] D. Roberts, M. Gardner, R. Church, S. Ustin, G. Scheer, and R. Green, "Mapping chaparral in the Santa Monica Mountains using multiple end-member spectral mixture models," *Remote Sens. Environ.*, vol. 65, no. 3, pp. 267–279, Sep. 1998.
- [103] Z. Huang, X. Jia, B. J. Turner, W. J. Foley, S. J. Dury, and I. R. Wallis, "Estimating the nitrogen concentration of eucalypt foliage by automatically extracting tree spectra from HYMAP data," *Photogramm. Eng. Remote Sens.*, vol. 73, no. 4, pp. 397–401, Apr. 2007.
- [104] B. Somers, S. Delalieux, J. Stuckens, W. W. Verstraeten, and P. Coppin, "A weighted linear spectral mixture analysis approach to address end-member variability in agricultural production systems," *Int. J. Remote Sens.*, vol. 30, no. 1, pp. 139–147, Jan. 2009.
- [105] R. N. Clark, G. A. Swayze, K. E. Livo, R. F. Kokaly, S. J. Sutley, J. B. Dalton, R. R. McDougal, and C. A. Gent, "Imaging spectroscopy: Earth and planetary remote sensing with the USGS tetracorder and expert systems," *J. Geophys. Res.*, vol. 108, no. 5131, pp. 1–44, Dec. 2003.
- [106] G. Asner and D. Lobell, "A biogeophysical approach for automated SWIR unmixing of soils and vegetation," *Remote Sens. Environ.*, vol. 74, no. 1, pp. 99–112, Oct. 2000.
- [107] G. P. Asner and K. B. Heidebrecht, "Spectral unmixing of vegetation, soil, and dry carbon cover in arid regions: Comparing multispectral and hyperspectral observations," *Int. J. Remote Sens.*, vol. 23, no. 19, pp. 3939–3958, Oct. 2002.
- [108] T. Rashed, J. R. Weeks, D. A. Roberts, J. Rogan, and R. Powell, "Measuring the physical composition of urban morphology using multiple end-member spectral mixture models," *Photogramm. Eng. Remote Sens.*, vol. 69, no. 9, pp. 1011–1020, Sep. 2003.
- [109] C. Wu, "Normalized spectral mixture analysis for monitoring urban composition using ETM+ imagery," *Remote Sens. Environ.*, vol. 93, no. 4, pp. 480–492, Dec. 2004.

- [110] C. Song, "Spectral mixture analysis for subpixel vegetation fractions in the urban environment: How to incorporate endmember variability?" *Remote Sens. Environ.*, vol. 95, no. 2, pp. 248–263, Mar. 2005.
- [111] S. Moussaoui, H. Hauksdóttir, F. Schmidt, C. Jutten, J. Chanussot, D. Brie, S. Doutó, and J. A. Benediktsson, "On the decomposition of Mars hyperspectral data by ICA and Bayesian positive source separation," *Neurocomputing*, vol. 71, no. 10–12, pp. 2194–2208, Jun. 2008.
- [112] M. Parente, J. L. Bishop, and J. F. Bell, "Spectral unmixing for mineral identification in pancam images of soils in Gusev crater, Mars," *Icarus*, vol. 203, no. 2, pp. 421–436, Oct. 2009.
- [113] M. Parente, J. T. Clark, A. J. Brown, and J. L. Bishop, "End-to-end simulation and analytical model of remote-sensing systems: Application to CRISM," *IEEE Trans. Geosci. Remote Sens.*, vol. 48, no. 11, pp. 3877–3888, Nov. 2010.
- [114] H. Du, W. Fan, G. Zhou, X. Xu, H. Ge, Y. Shi, Y. Zhou, R. Cui, and Y. Lü, "Retrieval of canopy closure and LAI of Moso bamboo forest using spectral mixture analysis based on real scenario simulation," *IEEE Trans. Geosci. Remote Sens.*, vol. 49, no. 11, pp. 4328–4340, Nov. 2011.
- [115] X. Ceamanos, S. Douté, B. Luo, F. Schmidt, G. Jouannic, and J. Chanussot, "Intercomparison and validation of techniques for spectral unmixing of hyperspectral images: A planetary case study," *IEEE Trans. Geoscience and Remote Sensing*, vol. 49, no. 11, pp. 4341–4358, Nov. 2011.
- [116] P. Addabbo, M. di Bisceglie, and C. Galdi, "The unmixing of atmospheric trace gases from hyperspectral satellite data," *IEEE Trans. Geosci. Remote Sens.*, vol. 50, no. 6, Jun. 2012, to be published.



Antonio Plaza (M'05–SM'07) received the M.S. and Ph.D. degrees in computer engineering from the University of Extremadura, Caceres, Spain.

He was a Visiting Researcher with the Remote Sensing Signal and Image Processing Laboratory, University of Maryland Baltimore County, Baltimore, with the Applied Information Sciences Branch, Goddard Space Flight Center, Greenbelt, MD, and with the Airborne Visible/Infrared Imaging Spectrometer Data Facility, Jet Propulsion Laboratory, Pasadena, CA. He is currently an Associate Professor with the Department of Technology of Computers and Communications, University of Extremadura, where he is the Head of the Hyperspectral Computing Laboratory. He was the Coordinator of the Hyperspectral Imaging Network, a European project designed to build an interdisciplinary research community focused on hyperspectral imaging activities. He has been a Proposal Reviewer with the European Commission, the European Space Agency, and the Spanish Government. He is the author or coauthor of around 300 publications on remotely sensed hyperspectral imaging, including more than 50 Journal Citation Report papers,

20 book chapters, and over 200 conference proceeding papers. His research interests include remotely sensed hyperspectral imaging, pattern recognition, signal and image processing, and efficient implementation of large-scale scientific problems on parallel and distributed computer architectures.

Dr. Plaza has coedited a book on high-performance computing in remote sensing and guest edited seven special issues on remotely sensed hyperspectral imaging for different journals, including the IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING (TGARS) (for which he has served as an Associate Editor on hyperspectral image analysis and signal processing since 2007), the IEEE JOURNAL OF SELECTED TOPICS IN APPLIED EARTH OBSERVATIONS AND REMOTE SENSING, the *International Journal of High Performance Computing Applications*, and the *Journal of Real-Time Image Processing*. He has served as a reviewer for more than 280 manuscripts submitted to more than 50 different journals, including more than 140 manuscripts reviewed for the IEEE TGARS. He served as a Chair for the IEEE Workshop on Hyperspectral Image and Signal Processing: Evolution in Remote Sensing in 2011. He has also been serving as a Chair for The International Society for Optics and Photonics Conference on Satellite Data Compression, Communications, and Processing since 2009 and for the SPIE Remote Sensing Europe Conference on High Performance Computing in Remote Sensing since 2011. He is currently serving as the Director of Education activities for the IEEE Geoscience and Remote Sensing Society. He was a recipient of the recognition of Best Reviewers of the IEEE GEOSCIENCE AND REMOTE SENSING LETTERS in 2009 and a recipient of the recognition of Best Reviewers of the IEEE TGARS in 2010.



Qian Du (S'98–M'00–SM'05) is currently an Associate Professor with the Department of Electrical and Computer Engineering, Mississippi State University, Mississippi State. She has authored or coauthored more than 160 scientific publications, including journal papers, book chapters, and peer-reviewed conference proceedings. Her research field is digital image processing and its application to remote sensing problems with an expertise on hyperspectral image exploitation. The research that she has conducted covers almost all the topics in remote sensing image processing and analysis, such as target detection, anomaly detection, change detection, supervised and unsupervised classification, linear and nonlinear unmixing, endmember extraction, real-time processing, parallel processing, band selection, data compression, registration and mosaicking, sharpening, visualization, etc. Her research interests also include image superresolution and neural networks.

Dr. Du has been active in the IEEE Geoscience and Remote Sensing Society (GRSS). She has been serving as a technical reviewer for many remote sensing and image processing journals and was the recipient of the 2010 Best Reviewer Award from IEEE GEOSCIENCE AND REMOTE SENSING LETTERS. She currently serves as the Cochair for the Data Fusion Technical Committee of IEEE GRSS. She serves as the Guest Editor for the special issue on Spectral Unmixing of Remotely Sensed Data for the IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING, for the special issue on High Performance Computing in Earth Observation and Remote Sensing, and for the special issue on Exploitation of Optical Multiangular Data for the IEEE JOURNAL OF SELECTED TOPICS IN APPLIED EARTH OBSERVATIONS AND REMOTE SENSING. She is the General Chair for the IEEE-GRSS 4th Workshop on Hyperspectral Image and Signal Processing: Evolution in Remote Sensing.



José M. Bioucas-Dias (S'87–M'95) received the E.E., M.Sc., Ph.D., and “Agregado” degrees in electrical and computer engineering from the Technical University of Lisbon, Lisbon, Portugal, in 1985, 1991, 1995, and 2007, respectively.

Since 1995, he has been with the Department of Electrical and Computer Engineering, Instituto Superior Tecnico. He is also a Senior Researcher with the Pattern and Image Analysis group at the Telecommunications Institute, which is a private nonprofit research institution. His research interests include signal and image processing, pattern recognition, optimization, and remote sensing. He was and is involved in several national and international research projects and networks, including the Marie Curie Actions “Hyperspectral Imaging Network” and the “European Doctoral Program in Signal Processing (SIGNAL).”

Dr. Bioucas-Dias is an Associate Editor of the IEEE TRANSACTIONS ON IMAGE PROCESSING, and he was an Associate Editor of the IEEE TRANSACTIONS ON CIRCUITS AND SYSTEMS. He is a Guest Editor of two IEEE special issues (IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING and IEEE JOURNAL OF SELECTED TOPICS IN APPLIED EARTH OBSERVATIONS AND REMOTE SENSING). He has been a member of program/technical committees of several international conferences. He was the General Cochair of the 3rd IEEE Workshop on Hyperspectral Image and Signal Processing: Evolution in Remote Sensing in 2011.



Xiuping Jia (M'93–SM'03) received the B.Eng. degree from the Beijing University of Posts and Telecommunications, Beijing, China, in 1982, and the Ph.D. degree in electrical engineering from The University of New South Wales, Sydney, Australia, in 1996.

She was a Visiting Scholar at several organizations during her sabbatical leaves, including Purdue University, West Lafayette, IN, Australian National University, Canberra, Australia, Beijing Normal University, Beijing, China, The Chinese University of Hong Kong, Shatin, Hong Kong, and The Chinese Academy of Sciences, China. She has given invited talks to the remote sensing research groups in several organizations, such as The Harbin Institute of Technology, Harbin, China, ITC, Netherlands, The Council for Scientific and Industrial Research in South Africa, Rochester Institute of Technology, Henrietta, NY, Center for Space and Remote Sensing Research, Taiwan, Shenzhen Institutes of Advanced Technology, China, and The University of Copenhagen, Copenhagen, Denmark. She delivered a plenary talk at the 2nd Workshop on Hyperspectral Image and Signal Processing, Iceland, in 2010. She is a Guest Professor of Harbin

Engineering University, Harbin. She chairs the Geoscience and Remote Sensing Society chapter of the Australian Capital Territory and New South Wales section, Australia. Since 1988, she has been with the School of Engineering and Information Technology, University College, The University of New South Wales, Australian Defence Force Academy, Canberra, Australia, where she is currently a Senior Lecturer. Her research areas are image processing, data analysis, and remote sensing applications. The projects that she has been involved in and supervised range from image registration, data compression, feature reduction, to spectral–spatial-based classification. Subpixel mapping has been addressed in recent years via spectral unmixing techniques and superresolution reconstruction approaches. She is the author or coauthor of over 100 publications on remote sensing image analysis, including journal papers, book chapters, and peer-reviewed conference proceedings. She is the coauthor of the remote sensing textbook entitled *Remote Sensing Digital Image Analysis* [Springer-Verlag, 3rd (1999) and 4th eds. (2006)].

Dr. Jia is an Associate Editor of the IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING.



Fred A. Kruse (M'96) received the B.S. degree in geology from the University of Massachusetts, Amherst, in 1976, and the M.S. and Ph.D. degrees in geology from the Colorado School of Mines, Golden, in 1984 and 1987, respectively.

He has been involved in remote sensing research and applications for over 28 years in positions with the U.S. Geological Survey, the University of Colorado, Boulder, the University of Nevada, Reno, and private industry. He is currently a Research Professor with the Department of Physics and the Remote Sensing Center, Naval Postgraduate School, Monterey, CA. He teaches courses in spectral remote sensing and synthetic aperture radar (SAR) and is developing Internet and computer-media-based remote sensing curricula. He is also one of the scientists that originally developed the image analysis software “ENVI.” His research focuses on physics-based exploration of diverse visible/near infrared, short-wave infrared, and long-wave infrared imaging spectrometer data and multifrequency, polarimetric, and interferometric SAR remote sensing data sets. Interests include the extraction of information applicable to scientific problems in

geology (geologic mapping, exploration for ore deposits, geothermal/hydrothermal systems, geologic hazards, and environmental monitoring) and for military and intelligence purposes.