

Received 23 January 2017; revised 14 August 2017; accepted 9 January 2018.  
Date of publication 29 March 2018; date of current version 4 April 2018.

Digital Object Identifier 10.1109/JTEHM.2018.2797983

# Height and Weight Estimation From Anthropometric Measurements Using Machine Learning Regressions

**DIEGO RATIVA<sup>ID</sup>, (Member, IEEE), BRUNO J. T. FERNANDES, (Member, IEEE),  
AND ALEXANDRE ROQUE**

Polytechnique School of Pernambuco, University of Pernambuco, Recife-Pernambuco 50720-001, Brazil

CORRESPONDING AUTHOR: D. RATIVA (diego.rativa@ecomppoli.br)

This work was supported by the Brazilian agencies, the Conselho Nacional de Desenvolvimento Científico e Tecnológico and the Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco.

**ABSTRACT** Height and weight are measurements explored to tracking nutritional diseases, energy expenditure, clinical conditions, drug dosages, and infusion rates. Many patients are not ambulant or may be unable to communicate, and a sequence of these factors may not allow accurate estimation or measurements; in those cases, it can be estimated approximately by anthropometric means. Different groups have proposed different linear or non-linear equations which coefficients are obtained by using single or multiple linear regressions. In this paper, we present a complete study of the application of different learning models to estimate height and weight from anthropometric measurements: support vector regression, Gaussian process, and artificial neural networks. The predicted values are significantly more accurate than that obtained with conventional linear regressions. In all the cases, the predictions are non-sensitive to ethnicity, and to gender, if more than two anthropometric parameters are analyzed. The learning model analysis creates new opportunities for anthropometric applications in industry, textile technology, security, and health care.

**INDEX TERMS** Machine learning, statistical learning, health information management.

## I. INTRODUCTION

Height and Weight are measurements broadly explored to tracking child's growth, nutritional diseases, energy expenditure, clinical conditions and health status [1]; Patients regularly are not ambulant or may be unable to communicate, and a sequence of these factors may not allow accurate estimation or measurements [2], [3].

Intensive care unit scoring systems, drug dosages, and infusion rates are commonly based on body weight, as well as, height is usually used to obtain a relation of healthy versus unhealthy weight [4]. Therefore, a correct estimation of height and weight of critically ill patients are critical for an adequate clinical care [1], [2].

W. C. Chumlea has been one of the first researchers to proposed linear equations to predict body height [1], [5], [6] and weight [1], [7] from anthropometric measurements for an elderly population. The equations are based on linear regressions, developed for a selected population in the USA, predicting weight within 95% confidence limits in

a range of 7.60-8.96kg, as well as, height with standard errors between 7.84cm and 8.44cm, depending on gender and Ethnicity; Therefore, the equations may be inappropriate for other populations [1].

Different groups have proposed subtly different weight/height prediction equations, based on single and multiple linear regressions, exploring other anthropometric parameters or analyzing other populations, achieving similar standard errors [8]–[12]. Linear regressions are attractive models because their representations are simple, and a straightforward algebra returns an analytical solution [13], that can be easily implemented by a healthcare professional.

Contemporaneously with the Chumlea's work, the Machine Learning (ML) field and its models from statistics and probability theory began to play an important role in research, engineering, economy, health, etc. [14]–[16]. ML is closely related to computational statistics, and it is defined as the development of algorithms that learn and make predictions from data or experience [17]. ML algorithms can

find patterns in complex scenarios, usually impossible to be identified by humans [18], therefore, ML regressions are usually more accurate than conventional regressions [13].

Recently, kernel machines have been presented as an appropriated approach for regression of biometric data [19]–[21]. As stated by Scholkopf, kernel machines provide modularity in the design, enabling easy combination with different learning algorithms, and have few parameters to be tuned in comparison with other models such as neural networks in an optimization procedure without spurious local minima [22], [23]. Moreover, other works have demonstrated that kernel machines have a strong founding theory, providing a more appropriated data representation for the problem that has been studied [24], [25].

Therefore, in this work, we aim to analyze the regression capability of kernel machines in comparison with traditional approaches for height and weight estimation from anthropometric measurements. We consider both sparse and non-sparse kernel machines - Support Vector Regression and Gaussian Process, respectively - to address the regression problem. Additionally, we also include a comparison with the results obtained by neural networks since such models have been extensively applied in machine learning problems recently. The method is demonstrated using the Third National Health and Nutrition Examination Survey [26] and the U.S. Army Anthropometry Survey [27] databases. Finally, the potentiality of the method is explored in a scenario of a patient without or partial mobility.

## II. METHODS

### A. STATISTICAL REGRESSIONS

In statistical regression, the relationships are modeled using linear or nonlinear predictor functions whose unknown model parameters are estimated from the experimental data. For instance, a mapping has two or more explanatory variables, given by [13]:

$$f_{\theta}(\mathbf{x}) = \langle \mathbf{w}, \mathbf{x} \rangle + b; \quad \theta = \{\mathbf{w}, b\} \quad (1)$$

where  $\langle \mathbf{w}, \mathbf{x} \rangle$  is the dot product between the weight vector  $\mathbf{w}$  and the inputs values  $\mathbf{x}$  and  $b$  is a bias or residual error, conventional regression models are often fitted using the least squares approach. For example, Chumlea *et al.* [5] establish a linear prediction equation where  $f_1$  is height,  $\mathbf{x}_1$  is the knee height, and  $\mathbf{w}_1$  is a multiplicative coefficient different for each analysed population.

Statistical regressions are attractive models to drawing conclusions about general principles based on a set of observations. In this work, the results obtained for different statistical traditional regressions are similar to the obtained with linear regressions (LR); for that reason, our discussion is limited to LR.


### B. MACHINE LEARNING REGRESSIONS

ML regression models are highly adaptable and are capable of modeling complex relationships; it can be implemented instead of statistical regression for the following cases:


the primary goal is to predict rather than explain, predictor variables are correlated or have non-linear relationships to the target variable, and there are numerous complex variable interactions.

As represented in Figure 1, in the case of ML Regression, instead of defined functions (Hand-crafted model) as used in statistical regression, one part of the existing data (i.e. Training and Validation Set) is used to identify settings for the model parameters that return the best realistic predictive performance, building and tuning the model. Another part of the data with new examples (i.e. Test Set) is used to test the performance of the regression model [28].

Traditional Regression:



Machine Learning Regression:



**FIGURE 1. Comparison between traditional and machine learning work-flow regressions.**

Sparse kernel machines are those one in which the kernel evaluation needs only a subset of the training data. In SVMs, for example, as less support vectors are used, sparser is the model since it will assign zero weights to most training patterns. Therefore, usually sparsity is a desired behavior because it gives a better interpretability and provides much faster solutions reducing the computational cost of the model. However, the information carried in the patterns with assigned zero weights is lost and it could be important in some type of problems [29], in special, those problems that are more susceptible to overfitting.

#### 1) SUPPORT VECTOR REGRESSION (SPARSE)

Support Vector Regression (SVR) with different Kernel functions such as Linear, Polynomial, and Gaussian function were explored. SVR with Gaussian Kernel achieved the best performance, the Gaussian base function is given by  $G(x_1, x_2) = \langle \phi(x_1), \phi(x_2) \rangle$ , where  $\phi(x)$  is a transformation that maps  $x$  to a high-dimensional space. For that reason, in this work our discussion is limited to SVR with Gaussian Kernel [30].

#### 2) GAUSSIAN PROCESSES REGRESSION (NON-SPARSE)

Gaussian process Regression (GPR) uses lazy learning methods and a similarity measure between points to predict values for unseen points from training data, more details can be

found in [31] and [32]. Therefore, it is possible to compute the prediction intervals using the trained model.

### 3) NEURAL NETWORK REGRESSION

The Neural Network Regression (NNR) is composed by Two-layer feed-forward with ten hidden sigmoid neurons. A Levenberg-Marquardt algorithm has been explored for training [33].

In this work, 70% of the analyzed data corresponds to the training group (train the ML regression and adjust the model weights); other 15% is the validation group, used to determine whether the classification accuracy of the ML regression is increasing or over-fitting. Finally, the rest 15% is going to test the trained ML model. The weight and height prediction performance of the different models were assessed using the Root Mean Square Error (RMSE). RMSE amplifies and severely punishes significant errors, and it is more appropriate to gauge model performance [34]. Moreover, it is on the same scale and units of the analyzed parameter (i.e. weight in kilograms and height in centimeters).

### C. DATABASES

#### 1) NHANES III

The Third National Health and Nutrition Examination Survey were collected between 1988 and 1994, contains data for 33,994 persons. Three ethnic groups are analyzed, non-Hispanic white (40%), non-Hispanic Black (30%) and Mexican-American (30%) subgroups for the US population [26]. In this work, we have selected adult subjects (over 21 years), with full data about anthropometric parameters related with height and weight corresponding to 4261 and 14783 subjects, respectively. Details about the measurement methodology to collect data can be found in [5] and [26].

#### 2) ANSUR


The 1988 U.S. Army Anthropometry Survey is one of the most widely used anthropometry databases because of the rigorous methodology and a large number of measures [27]. The database is composed of two Gender subgroups: Male (1774) and Female (2208) subjects, totalizing 3982 samples.

### III. ANTHROPOMETRIC CORRELATION ANALYSIS


There are several correlations between anthropometric measurements of the human body, such as body height and Member-height, as well as, weight and Member-circumference. In Figure 2 are represented, usual correlations explored in healthcare: Height vs. Knee-height (a) and weight vs. Buttock-circumference (b).

The most representative anthropometric Square Correlation Coefficients ( $R^2$ ) for weight and height are summarized in Figure 3. The correlation analysis has been elaborated for a non-Hispanic white male group (NHANES III, 1369 subjects) and a Male Group (ANSUR, 1776 subjects).

Weight is often related with Waist-circumference [5], without regard that there are high correlations with Buttock, Arm, Calf, Biceps, and Chest circumference as well. Therefore,



**FIGURE 2.** Scatter diagrams for a non-Hispanic white male group (1369 subjects): Height vs. Knee-height (a) and weight vs. Buttock-circumference (b).




**FIGURE 3.** Weight and Member-circumference (left), as well as, height and Member-height (right) correlations for a non-Hispanic white Male Group (NHANES III, 1369 subjects) and a Male Group (ANSUR, 1776 subjects).

an individual or combinational parameter analysis of other parameters may be explored in cases where is not possible to measure Waist-circumference.

An analysis of the predicted weight and height for different individual or combinations of Anthropometric Measurements is illustrated in the following subsections.

#### A. PREDICTED WEIGHT

Three cases of the multiple possible combinations of anthropometric parameters are explored to analyze the performance to predict weight of the ML regressions. Anthropometric data corresponds to 14783 adult subjects from NHANES III database.



**FIGURE 4.** Predicted vs Measured weight scatter diagram; for Linear (L), Support Vector (S), Gaussian Process (G) and Neural Network (N) regressions. Anthropometric parameters explored are: (a) Waist circumference (Waist); (b) Waist, Height, Buttock circumference (Butt.), Thigh Circumference (Thigh), Gender, Ethnicity (Ethn.); (c) Waist, Height, Butt., Thigh, Arm Circumference (Arm), Gender and Ethn. Group of 14783 adult subjects from NHANES III.

The combination of anthropometric parameter inputs are organized as follow:

- **Comb. 1:** Waist circumference;
- **Comb. 2:** Gender, Ethnicity, height, as well as, Buttock and Waist circumferences;
- **Comb. 3:** Comb. 2 in addition to Arm circumference.


Figure 4 shows the scatter diagram of the Predicted vs. Measured weight of the proposed combinations obtained by Linear and ML regressions; a solid line is used to represent a perfect correlation ( $R^2 = 1$ ).

For Comb. 1, (Fig. 4(a)) where the Waist circumference parameter is the only input parameter, the RMSE values obtained are similar to the reported in the literature [5], which confidence interval (CI) is next to 78% and limits between 16.55kg and 18.49kg.

Furthermore, in the cases of Comb 2. and Comb. 3 (Fig. 5(b) and (c)), if more parameters are explored to train the models, the prediction is much more accurate. A CI of 95% with limits of 3.20kg and 3.50kg is obtained with linear regression, whereas, the best prediction performance is achieved with GPR, CI of 98% with limits of 1.80kg and 2.25kg. The prediction performance of NNR and SVR are subtly larger than the obtained with GPR.

Unlike ML regressions, the weight prediction with conventional regressions fails ( $c_L$ ) for values smaller than 50kg and higher than 110kg, as shown in Figure 4. (c). The same behavior is observed for other possible combinations of anthropometric values. In Figure 5, it can be seen the tendency of the weight prediction with a LR and its difference with a perfect correlation (Solid Line).

Examples of the prediction model performance for different anthropometric parameter combinations are summarized



**FIGURE 5.** Scatter diagram of the Predicted vs. Measured weight by using a conventional (Linear) regression, for a group of adult subjects from the NHANES III database (14783 subjects); (a) lighter 50kg (approx. 1500 subjects), and (b) heavier than 110kg (approx. 300 subjects). The solid line is a perfect weight prediction.

in Table 1; each subtable is organized by the number of anthropometric parameters explored as input: one, two (vertical and a horizontal parameter) and four parameters.

The ML regressions performance is better than conventional regression, for all possible combinations. RMSE values for the predicted weight obtained from SVR and GPR are similars (difference in the third decimal place); however, the accuracy in terms of CI is subtly better for GPR.

## B. PREDICTED HEIGHT

The correlation between predicted and measured height is analyzed for individual or different combinations of



**TABLE 1.** Weight prediction performance (RMSE in kilograms) of linear, support vector, gaussian process, and neural network regressions; for 14783 adult subjects (NHANES III). The anthropometric parameters input are height, gender, Ethnicity (Ethn.), as well as, Buttock (Butt.), Thigh (Thigh), Waist, (Waist) and Arm circumferences (Arm).

Anthropom. Parameter		Linear Regression										
		One Param.	Two Param.						Four Param. + Gender + Ethn.			
			+Height	+Butt.	+Thigh	+Waist	+Arm	+Gender	+Ethn.	+Height +Arm +Butt.	+Height +Arm +Thigh	+Height +Arm +Waist
Anthropom. Parameter	Height	14.30	–	6.69	8.33	6.81	6.48	14.30	14.30	–	4.32	3.70
	Butt.	9.95	6.69	–	9.04	8.10	7.33	6.38	9.94	–	–	–
	Thigh	10.00	8.33	9.04	–	6.59	7.54	8.43	10.00	4.32	–	3.60
	Waist	9.06	6.48	8.10	6.59	–	6.90	8.54	9.04	3.70	3.60	–
	Arm	8.11	6.48	7.33	7.54	6.90	–	7.66	8.00	–	–	–

Anthropom. Parameter		Support Vector Regression/Gaussian Process Regression										
		One Param.	Two Param.						Four Param. + Gender + Ethn.			
			+Height	+Butt.	+Thigh	+Waist	+Arm	+Gender	+Ethn.	+Height +Arm +Butt.	+Height +Arm +Thigh	+Height +Arm +Waist
Anthropom. Parameter	Height	7.07	–	5.88	6.75	5.72	5.35	7.02	7.10	–	3.60	3.00
	Butt.	7.14	5.88	–	7.02	6.95	6.52	5.63	7.00	–	–	–
	Thigh	6.06	6.75	7.02	–	5.94	6.58	6.05	6.06	3.60	–	2.97
	Waist	7.16	5.72	6.95	5.94	–	5.86	7.08	7.15	3.00	2.97	–
	Arm	7.10	5.35	6.52	6.58	5.86	–	6.62	6.91	–	–	–

Anthropom. Parameter		Neural Network Regression										
		One Param.	Two Param.						Four Param. + Gender + Ethn.			
			+Height	+Butt.	+Thigh	+Waist	+Arm	+Gender	+Ethn.	+Height +Arm +Butt.	+Height +Arm +Thigh	+Height +Arm +Waist
Anthropom. Parameter	Height	7.11	–	5.81	6.68	5.77	5.37	7.11	7.08	–	3.71	3.14
	Butt.	7.42	5.81	–	7.13	6.83	6.46	5.64	7.41	–	–	–
	Thigh	5.99	6.68	7.13	–	5.93	6.62	5.98	5.99	3.71	–	3.00
	Waist	7.50	5.77	6.83	5.93	–	5.83	7.18	7.31	3.14	3.04	–
	Arm	7.03	5.37	6.46	6.62	5.83	–	6.65	6.94	–	–	–

**TABLE 2.** Height prediction performance (RMSE in centimeters) of linear, support vector, gaussian process, and neural network regressions; for 4261 adult subjects (NHANES III). The anthropometric parameters input are Knee height (Knee), Gender, Ethnicity (Ethn.), Functional leg (Leg) and forearm (Arm) lengths, as well as, Wrist (Wrist) and sitting (Sit) heights.


Anthropom. Parameter		Linear Regression										
		One Param.	Two Param.						Four Param. + Gender + Ethn.			
			+Knee	+Leg	+Arm	+Wrist	+Sit.	+Gender	+Ethn.	+Knee +Leg +Arm	+Knee +Leg +Wrist	+Knee +Leg +Wrist
Anthropom. Parameter	Knee	4.85	–	4.57	4.61	4.60	2.43	4.61	4.08	–	–	–
	Leg	7.10	4.57	–	5.48	5.77	3.58	4.75	7.06	–	–	–
	Arm	6.29	4.61	5.48	–	5.67	3.55	5.42	6.24	–	4.03	2.20
	Wrist	7.01	4.60	5.77	5.67	–	4.00	6.43	6.89	3.93	–	2.20
	Sit.	4.45	2.43	3.58	3.55	4.00	–	4.18	4.43	2.17	2.20	–

Anthropom. Parameter		Support Vector Regression/Gaussian Process Regression										
		One Param.	Two Param.						Four Param. + Gender + Ethn.			
			+Knee	+Leg	+Arm	+Wrist	+Sit.	+Gender	+Ethn.	+Knee +Leg +Arm	+Knee +Leg +Wrist	+Knee +Leg +Wrist
Anthropom. Parameter	Knee	4.05	–	3.86	3.83	3.90	2.12	3.62	4.01	–	–	–
	Leg	5.18	3.86	–	4.58	4.59	3.29	4.65	5.12	–	–	–
	Arm	4.94	3.83	4.58	–	4.53	3.26	4.48	4.80	–	3.34	1.89
	Wrist	4.86	3.90	4.59	4.53	–	3.36	4.80	4.93	3.26	–	1.86
	Sit.	3.98	2.12	3.29	3.26	3.36	–	3.77	3.64	1.86	1.86	–

Anthropom. Parameter		Neural Network Regression										
		One Param.	Two Param.						Four Param. + Gender + Ethn.			
			+Knee	+Leg	+Arm	+Wrist	+Sit.	+Gender	+Ethn.	+Knee +Leg +Arm	+Knee +Leg +Wrist	+Knee +Leg +Wrist
Anthropom. Parameter	Knee	4.03	–	3.83	3.84	3.80	2.11	3.69	4.08	–	–	–
	Leg	4.89	3.83	–	4.51	4.57	3.29	4.62	5.03	–	–	–
	Arm	4.80	3.83	4.51	–	4.56	3.29	4.40	4.62	–	3.44	1.96
	Wrist	4.88	3.80	4.57	4.56	–	3.39	4.88	4.87	3.42	–	1.91
	Sit.	3.93	2.11	3.29	3.29	3.39	–	3.76	3.65	1.91	1.94	–



**FIGURE 6.** RMSE values for weight (a) and height prediction (b) for Male and Female groups analyzed jointly using a GPR, where Gender and Non-Gender means using or not the Gender as input parameter, respectively. Group of 14783 adult subjects from NHANES III.

anthropometric parameter inputs, parameters with more correlation with body's height have been selected for the analysis.


Thus, the anthropometric parameters entries are Knee-height, Gender, Ethnicity, functional-Leg and Forearm-hand lengths, as well as, Wrist and Sitting heights. The Height prediction performance values (RMSE in centimeters) of the explored models are summarized in Table 2. Similarly to the weight prediction case, the performance of the ML regressions is better than conventional regression, for all possible combinations.

A CI of 94% with limits of 11.2cm and 13.5cm is obtained with conventional regression models, whereas, a CI of 95% with limits of 6.7cm and 9.0cm, is achieved with GPR.


For comparison purposes, the Weight and Height prediction performance (RMSE and R-square values) obtained with the proposed model, as well as, the most representative values previously reported in the literature are summarized in Table 3. Our model reaches an RMSE value of 2.11 and an R-Squared of 0.84, for Weight and Height predictions, respectively. Therefore, for both predictions, the GPR method has a better performance than the linear methods.

### C. GENDER AND ETHNICITY PARAMETERS

It is well-known that anthropometric correlation values are different for male and female subjects, such that weight and height estimation equations have different coefficients for each gender. Furthermore, different equations have been reported for groups in various continents [5], [6]. In our analysis, all subjects are analyzed together, and Gender and Ethnicity are input parameters of the ML regressions.



**FIGURE 7.** True success rate of a support vector machine to classify gender, parameters explored for weight (a) and height (b) estimations. Group of 14783 adult subjects from NHANES III.



**FIGURE 8.** RMSE values for weight (a) and height prediction (b), where Ethnicity and Non-Ethnicity means using or not the Ethnicity as input parameter, respectively. Group of 14783 adult subjects from NHANES III.

#### 1) GENDER ANALYSIS

As can be observed in Tables 1 and 2, weight and height estimations are more accurate if the Gender parameter (1 for male and 0 for female subjects) is considered. For example, in the case that Buttock circumference is the only one input, the weight prediction SVR model has an RMSE of 7.14kg. However, if the Gender parameter is a second input, the RMSE is reduced to 5.88kg.

Moreover, if more than two parameters are analyzed (e.g. Buttock and Waist circumferences), the prediction is not more sensitive to Gender, as shown in Figure 6.


**TABLE 3. Height and weight predictions performance from anthropometric measurements.**

Height	Author, year	n	Age (years)	Gender	Regression	Predictor variable	RMSE
	Chumlea et al, [7]	4,750	Over 60	Men/Women*	Linear	Sitting height, Age	3.25
Cereda et al., [35]	635	30-55	Men/Women*	Linear	Knee height, Tibia length, body composition	2.2 ** / 4.4***	
Proposed Model	4,261	Over 21	Men/Women	Gaussian Process	Sitting height	2.11	

Weight	Author, year	n	Age (years)	Gender	Regression	Predictor variable	R-Squared
	Lin, et al. [36]	235	Over 18	Men/Women*	Linear	Knee height, Mid Arm circumference	0.69
Proposed Model	4,261	Over 21	Men/Women	Gaussian Process	Knee height, Mid Arm circumference, Stature	0.84	

(\*separated groups, \*\*men, \*\*\*women)



Case (1). Lower Limb	Case (2). Upper Limb	Case (3). Depth Parameters	Case (4)	Case (5)	Case (6)
Gender (1) Knee height (2) Calf circ. (3) Knee circ. (4) Thigh circ. Distal (5) Thigh circ. Proximal	Gender (I) Acroradial Length (II) Axillary Arm circ. (III) Scye circ. (IV) Biceps circ. (V) Elbow circ.	Gender (I) Acroradial Length (D1) Chest Depth (D2) Waist Depth (D3) Buttock Depth	Lower and Upper Limbs	Upper Limb and Depth Param.	Lower and Upper Limbs and Depth Param.
<b>Confidence</b> 93% 4.05kg <b>Interval</b> 94% 5.16kg	90% 5.60kg 92% 6.88kg	90% 5.64kg 92% 6.92kg	96% 1.91kg 97% 2.70kg	94% 2.94kg 96% 3.91kg	97% 1.14kg 98% 1.77kg
<b>Case (1) + Height</b> <b>Confidence</b> 94% 3.19kg <b>Interval</b> 95% 4.20kg	<b>Case (2) + Height</b> 92% 4.61kg 93% 5.80kg	<b>Case (3) + Height</b> 93% 3.93kg 94% 5.04kg	<b>Case (4) + Height</b> 97% 1.48kg 98% 2.20kg	<b>Case (5) + Height</b> 96% 1.96kg 97% 2.77kg	<b>Case (6) + Height</b> 98% 0.64kg 99% 1.13kg

**FIGURE 9. Weight estimation from Lower and Upper limb, and Depth anthropometric parameters, that can be measured comfortably for patients without or partial mobility. Analysis based on the ANSUR database (3997 subjects).**

The effect can be explained by the property of the ML regression to identify the subject Gender. To test our hypothesis, a Support Vector Machine (SVM) is explored as a binary classifier to determine the subject Gender, using the same input parameters employed in the SVR. As can be seen in Figure 7, considering two input parameters, the SVM classifier has a true success rate (TSR) of 80% to classify the subjects Gender; exploring more than two parameters, the TRS reach values next to 90%.

Therefore, the Gender parameter plays an important role in the prediction if just one anthropometric parameter is explored as an input parameter, but it is redundant in the case that more than two parameters are analyzed by the ML regression.

**2) ETHNICITY NON-SENSITIVE**

As aforementioned, the NHANES III database is composed of three ethnic groups [26]. Therefore in our analysis, the parameters are established as non-Hispanic white (1), non-Hispanic Black (2), and Mexican-American (3) parameters.

The prediction performance has not an important dependence with Ethnicity, which RMSE values are similar in the

cases of the Ethnicity parameter is considered or not, as represented in Figure 8 and summarized in Table 1 and 2. The same non-Sensitivity to Ethnicity is found for the ANSUR database.

On the other hand, analogously to the Gender classification, an SVM was explored unsuccessfully to classify the Ethnicity of the subjects (TSR next to 50%).

**3) CROSS-VALIDATION DATABASES**

Cross-validation tests between the NHANES III and the ANSUR databases test have been implemented. As summarized in Table 4, RMSE values achieved with both Inter- and Intra- database analysis are quite similar, proving the generalization of the ML regression model.

**IV. ESTIMATING WEIGHT FOR A SUBJECT WITHOUT MOBILITY**

In this numeral, groups of anthropometric measurements are proposed to estimate weight based on anatomical regions exploring a Gaussian Process Regression. Therefore, the healthcare professional has different options to collect circumferences and member lengths; that can be measured comfortably for patients without or partial mobility. The

**TABLE 4. Cross-validation between the NHANESIII and ANSUR databases.**

Training Database	Test Database	RMSE (kg)
NHANES III	NHANES III	2.97
NHANES III	ANSUR	2.99
ANSUR	ANSUR	2.89
ANSUR	NHANES III	2.97

group of measurements are organized as follow:

- **Case 1. Lower Limb:** Knee-height, and Calf, Knee, Thigh Distal, Thigh Proximal circumferences;
- **Case 2. Upper Limb:** Acroradial-Length, and Axillary arm, Scye, Biceps flexed, Elbow circumferences;
- **Case 3. Depth parameters:** Acroradial-Length, and Chest, Waist and Buttock depths;
- **Case 4. Lower and Upper Limb;**
- **Case 5. Limbs and Depth parameters.**

Weight estimations are based on the ANSUR database, which contains anthropometric parameters such as Calf, Axillary Arm, and Biceps circumferences, as well as, Chest, Waist and Buttock depth. Male and Females groups are analyzed jointly, and Gender is an input parameter.

It is important to note that the ANSUR anthropometric measurements were obtained in a stand-up or sitting positions. Consequently, the values are subtly different of the obtained from a patient placed in a Recumbent or Fowler positions.

As represented in Figure 9, the weight prediction from the Lower limb has the best performance, with a CI of 94% and limits between 4.05 and 5.16kg. If the patients' height is a known parameter, the CI reaches values close to 95% and limits between 3.19 and 4.20kg.

In case that is possible to measure both limbs (lower and upper), the CI obtained is 97% with limits between 2.94 and 3.91kg. Furthermore, if Depth parameters can be measured as well, the prediction reach the best accuracy value with a CI of 98% and limits between 1.14 and 1.77kg, and 99% with limits between 0.64kg and 1.13kg if the Height subject is known.

## V. CONCLUSION

Weight and Height prediction performance from anthropometric measurements obtained with Machine Learning regression models is better than the achieved with conventional statistical regressions.

In particular, the non-sparse kernel machine (Gaussian Process Regression) shows the best performance. Despite having a more expensive computational cost, the Gaussian Process Regression has the ability to choose the kernel hyper parameters directly from the training data what makes it a more flexible model with fewer setup requirements.

Besides, we believe that the non-sparsity of the Gaussian Process Regression enabled the usage of important information that might be discarded in the SVM through the assignment of zero weight for some inputs.

Weight and Height can be predicted for Female and Male subjects with the same ML regression, where Gender is one

input parameter. On the other hand, the Weight and Height predictions obtained with ML regressions have not a dependence with ethnicity for both populations analyzed.

According to the Cross-validation database analysis, the ML models are robust and general for analysis of different populations. Trained models obtained with ML regressions can be embedded in applications software to run on mobile devices and smartphones, opening new opportunities for anthropometric applications in industry, textile technology, security, and health-care.

## REFERENCES

- [1] World Health Organization and others, "Physical status: The use and interpretation of anthropometry. Geneva; 1995," *WHO Tech. Rep. Ser.*, vol. 854, pp. 2009–2006, 2011.
- [2] R. Bloomfield, E. Steel, G. MacLennan, and D. W. Noble, "Accuracy of weight and height estimation in an intensive care unit: Implications for clinical practice and research," *Critical Care Med.*, vol. 34, no. 8, pp. 2153–2157, 2006.
- [3] B. L. Anglemeyer, C. Hernandez, J. H. Brice, and B. Zou, "The accuracy of visual estimation of body weight in the ED," *Amer. J. Emergency Med.*, vol. 22, no. 7, pp. 526–529, Nov. 2004.
- [4] K. B. Michels, S. Greenland, and B. A. Rosner, "Does body mass index adequately capture the relation of body composition and body size to health outcomes?" *Amer. J. Epidemiol.*, vol. 147, no. 2, pp. 167–172, 1998.
- [5] W. C. Chumlea, S. Guo, A. F. Roche, and M. L. Steinbaugh, "Prediction of body weight for the nonambulatory elderly from anthropometry," *J. Amer. Dietetic Assoc.*, vol. 88, no. 5, pp. 564–568, 1988.
- [6] W. C. Chumlea and S. Guo, "Equations for predicting stature in white and black elderly individuals," *J. Gerontol.*, vol. 47, no. 6, pp. M197–M203, 1992.
- [7] W. M. C. Chumlea, S. S. Guo, and M. L. Steinbaugh, "Prediction of stature from knee height for black and white adults and children with application to mobility-impaired or handicapped persons," *J. Amer. Dietetic Assoc.*, vol. 94, no. 12, pp. 1385–1391, 1994.
- [8] J. W. Prothro and C. A. Rosenbloom, "Physical measurements in an elderly black population: Knee height as the dominant indicator of stature," *J. Gerontol.*, vol. 48, no. 1, pp. M15–M18, 1993.
- [9] L. M. Gauld, J. Kappers, J. B. Carlin, and C. F. Robertson, "Height prediction from ulna length," *Develop. Med. Child Neurol.*, vol. 46, no. 7, pp. 475–480, 2004.
- [10] C. O. Mitchell and D. A. Lipschitz, "Arm length measurement as an alternative to height in nutritional assessment of the elderly," *J. Parenteral Enteral Nutrition*, vol. 6, no. 3, pp. 226–229, 1982.
- [11] B. Kuiti and K. Bose, "Predictive equations for height estimation using knee height of older Bengalees of Purba Medinipur, West Bengal, India," *Anthropol. Rev.*, vol. 79, no. 1, pp. 47–57, 2016.
- [12] S. E. Hogan, "Knee height as a predictor of recumbent length for individuals with mobility-impaired cerebral palsy," *J. Amer. College Nutrition*, vol. 18, no. 2, pp. 201–205, 1999.
- [13] D. A. Freedman, *Statistical Models: Theory and Practice*. Cambridge, U.K.: Cambridge Univ. Press, 2009.
- [14] J. H. Friedman, "Data mining and statistics: What is the connection?" *Comput. Sci. Statist.*, vol. 29, no. 1, pp. 3–9, 1998.
- [15] M. C. Grassi, A. M. Caricati, M. Intraligi, M. Buscema, and P. Nencini, "Artificial neural network assessment of substitutive pharmacological treatments in hospitalised intravenous drug users," *Artif. Intell. Med.*, vol. 24, no. 1, pp. 37–49, Jan. 2002.
- [16] P. J. Lisboa and A. F. G. Taktak, "The use of artificial neural networks in decision support in cancer: A systematic review," *Neural Netw.*, vol. 19, no. 4, pp. 408–415, May 2006.
- [17] C. M. Bishop, *Pattern Recognition and Machine Learning*. New York, NY, USA: Springer-Verlag, 2016.
- [18] V. Vapnik, *The Nature of Statistical Learning Theory*. New York, NY, USA: Springer, 2013.
- [19] T. Shepherd and R. Owenius, "Gaussian process models of dynamic PET for functional volume definition in radiation oncology," *IEEE Trans. Med. Imag.*, vol. 31, no. 8, pp. 1542–1556, Aug. 2012.



- [20] Y. Yun, H.-C. Kim, S. Y. Shin, J. Lee, A. D. Deshpande, and C. Kim, "Statistical method for prediction of gait kinematics with Gaussian process regression," *J. Biomech.*, vol. 47, no. 1, pp. 186–192, Jan. 2014.
- [21] F. Javed *et al.*, "RBF kernel based support vector regression to estimate the blood volume and heart rate responses during hemodialysis," in *Proc. Annu. Int. Conf. IEEE Eng. Med. Biol. Soc. (EMBC)*, Sep. 2009, pp. 4352–4355.
- [22] B. Schölkopf, C. J. C. Burges, and A. J. Smola, *Advances in Kernel Methods—Support Vector Learning*. Cambridge, MA, USA: MIT Press, 1999.
- [23] J. Shawe-Taylor and S. Sun, "Kernel methods and support vector machines," in *Academic Press Library in Signal Processing*, vol. 1. New York, NY, USA: Elsevier, 2014. pp. 857–881.
- [24] T. Hofmann, B. Schölkopf, and A. J. Smola, "Kernel methods in machine learning," *Ann. Statist.*, vol. 36, no. 3, pp. 1171–1220, 2008.
- [25] J. Shawe-Taylor and N. Cristianini, *Kernel Methods for Pattern Analysis*. Cambridge, U.K.: Cambridge Univ. Press, 2004.
- [26] *Third National Health and Nutrition Examination Survey, 1988–1994, NHANES III*, Nat. Center Health Statist., Hyattsville, MD, USA, 1996.
- [27] C. C. Gordon, T. Churchill, C. E. Clauser, B. Bradtmiller, and J. T. McConville, "Anthropometric survey of us army personnel: Methods and summary statistics 1988," DTIC Document, U.S. Army Natick Soldier Res., Develop. Eng. Center Natick, Natick, MA, USA, Tech. Rep., 1989.
- [28] M. Kuhn and K. Johnson, *Applied Predictive Modeling*. New York, NY, USA: Springer, 2013.
- [29] F. Yan, J. Kittler, K. Mikolajczyk, and A. Tahir, "Non-sparse multiple kernel Fisher discriminant analysis," *J. Mach. Learn. Res.*, vol. 13, no. 1, pp. 607–642, Mar. 2012.
- [30] O. Chapelle, V. Vapnik, O. Bousquet, and S. Mukherjee, "Choosing multiple parameters for support vector machines," *Mach. Learn.*, vol. 46, nos. 1–3, pp. 131–159, 2002.
- [31] C. E. Rasmussen, *Gaussian Processes for Machine Learning*. Cambridge, MA, USA: MIT Press, 2006.
- [32] M. Seeger, "Gaussian processes for machine learning," *Int. J. Neural Syst.*, vol. 14, no. 2, pp. 69–106, 2004.
- [33] B. D. Ripley, *Pattern Recognition and Neural Networks*. Cambridge, U.K.: Cambridge Univ. Press, 2007.
- [34] T. Chai and R. R. Draxler, "Root mean square error (RMSE) or mean absolute error (MAE)?—Arguments against avoiding RMSE in the literature," *Geosci. Model Develop.*, vol. 7, no. 3, pp. 1247–1250, 2014.
- [35] E. Cereda, S. Bertoli, and A. Battezzati, "Height prediction formula for middle-aged (30–55 y) Caucasians," *Nutrition*, vol. 26, nos. 11–12, pp. 1075–1081, Nov./Dec. 2010.
- [36] B. W. Lin, D. Yoshida, J. Quinn, and M. Strehlow, "A better way to estimate adult patients' weights," *Amer. J. Emergency Med.*, vol. 27, no. 9, pp. 1060–1064, 2009.